

# Permissive hypercapnia in acute respiratory distress syndrome — is now the time to get strict?

Alistair D Nichol

Although mechanical ventilation is frequently a life-saving intervention for patients with acute respiratory distress syndrome (ARDS), there is an increasing awareness that mechanical ventilation itself can initiate or exacerbate lung injury. Over the past two decades, there has been a major shift in how we ventilate these patients. Despite the increase in our understanding, the ideal ventilation strategy and the appropriate management of ventilation (control of  $P_{aCO_2}$ ) and oxygenation (control of  $P_{aO_2}$ ) for patients with ARDS is still controversial. Many clinicians still aim for physiologically “normal” arterial blood gas parameters and, in this pursuit, may induce additional pulmonary injury, thus increasing morbidity and mortality.

## Permissive hypercapnia — what’s the story?

The term “permissive hypercapnia” was coined after two case series by Hickling and colleagues in the early 1990s that suggested that limitation of airway pressure ( $P_{AW}$ ) and tidal volume ( $V_T$ ), with a tolerant approach to elevations in alveolar and arterial  $CO_2$  (permissive hypercapnia, mean maximum  $P_{aCO_2}$  66.5 mmHg)<sup>1</sup> was associated with lower hospital mortality than predicted by Acute Physiological and Chronic Health Evaluation (APACHE) II scores.<sup>1,2</sup> Subsequently, Amato and colleagues randomly allocated 53 patients with ARDS to receive either “protective” low  $V_T$  (<6 mL/kg), high positive-end expiratory pressure (PEEP) and a permissive approach to  $CO_2$  control (initial  $P_{aCO_2}$  up to 80 mmHg), or conventional  $V_T$  (12 mL/kg) with respiratory rate adjusted to maintain  $P_{aCO_2}$  in the normal range.<sup>3</sup> The protective ventilation strategy was associated with an increase in survival at 28 days. These results demonstrated that a protective ventilation strategy, which limited  $P_{AW}$  and  $V_T$  by tolerating increases in  $P_{aCO_2}$ , could be associated with a survival benefit among patients with ARDS.

However, the subsequent multicentre ARDSNet trial, which also demonstrated a survival advantage for a low  $V_T$  and  $P_{AW}$  strategy compared with the control group, used an active approach to prevent hypercapnia or acidosis (by increasing respiratory rate and  $V_T$  and administering sodium bicarbonate).<sup>4</sup> This suggested to some that it was irrelevant whether a permissive or active approach to the management of  $CO_2$  was adopted, as long as  $P_{AW}$  and  $V_T$  were limited.

However, there is a growing body of evidence suggesting that such an active approach may be deleterious, and that a

permissive approach minimises lung injury and facilitates the goals of the ideal protective ventilation strategies for patients with ARDS.

## Active control of carbon dioxide may be deleterious in acute respiratory distress syndrome

There appears to be no safe  $V_T$  or  $P_{AW}$  cut-off in ARDS, and there are additive advantages to further decreasing both, thus minimising volutrauma and barotrauma.<sup>5</sup> Furthermore, the cyclic opening and closing of acetalveolar units with each breath is deleterious.<sup>6</sup> The simple act of increasing the respiratory rate from 12 to 30 breaths per minute (to control  $CO_2$ ) adds over 25 000 additional opening and closing cycles per day to an already injured lung. These mechanical interactions also act as an engine for a systemic inflammatory response, multiorgan failure and death in patients with ARDS.<sup>7,8</sup> Therefore, increasing the  $V_T$  or respiratory rate to combat rises in  $CO_2$  may further augment lung injury and worsen outcomes. In addition, the exogenous administration of sodium bicarbonate to control pH may be ineffective and actually increase acidosis in ARDS patients, as well as being associated with other detrimental effects (ie, hypocalcaemia or hypokalaemia).<sup>9,10</sup> These findings suggest that the ARDSNet practice of increasing  $V_T$ , increasing respiratory rate, and infusing exogenous bicarbonate to control  $CO_2$  or acidosis may actually increase injury.

## Permissive hypercapnia may do more than allow low distension ventilation

Laboratory studies suggest that hypercapnic acidosis may have anti-inflammatory effects that can attenuate acute lung injury.<sup>11</sup> Fortunately, these immunomodulatory effects appear not to be deleterious in infective models of lung injury when appropriate antibiotic therapy is administered.<sup>12,13</sup> These findings suggest that hypercapnia is not just an associated side effect of ventilation strategies that aim to minimise lung stretch, but that it could have specific beneficial effects. Interestingly, a recent multivariate logistic regression analysis of the ARDSNet trial, after controlling for comorbidities and severity of lung injury, found that the patients who had hypercapnic acidosis on study Day 1 had a reduced 28-day mortality in the higher  $V_T$  group, a result consistent with a protective effect of hypercapnic acidosis

against ventilator-induced lung injury.<sup>14</sup> Taken together, these findings suggest that there are potential additional beneficial effects of a permissive approach to hypercapnia that may have been lost in the ARDSNet strategy.

### Permissive hypercapnia — time to be strict about its application in ARDS

Several low  $V_T$  prospective multicentre randomised controlled trials have been conducted in patients with acute lung injury or ARDS to compare high and low PEEP ( $\pm$ recruitment ventilation strategies).<sup>15,16</sup> While these trials demonstrated a clear improvement in oxygenation and a suggestion that high PEEP may reduce ventilator days, no survival benefit was noted in the high-PEEP groups. Despite these results, the interest in high-PEEP strategies has not diminished.<sup>17</sup> This is partly because of perceived design flaws of these trials; in particular, allowing similar ventilator driving pressures in the high-PEEP and the low-PEEP arms, resulting in higher plateau pressures in the high-PEEP arm of the trials, a clear confounding factor. This has prompted the development of newer strategies that not only aim to open the closed acetalatic segments with recruitment manoeuvres and higher PEEP, but also recognise that the use of ultralow tidal volumes are necessary in the face of higher PEEP to minimise airway pressure and realise the true potential benefits of protective open lung ventilator strategies.<sup>18</sup> To safely achieve these ventilator goals a permissive approach to hypercapnia is essential. Pilot work suggests that such an approach may be beneficial in ARDS.<sup>17</sup>

There is growing evidence in the management of critical illness that the short-term aim of “normal” physiological variables, although attractive, can actually result in worse long-term outcomes.<sup>19</sup> The time has come where we need to provide the most protective mechanical ventilation strategy if we are not adopting a deliberate policy of permissive hypercapnia for patients with ARDS.

### Author details

Alistair D Nichol, Associate Professor,<sup>1</sup> and Intensivist<sup>2</sup>

1 Australian and New Zealand Intensive Care Research Centre, School of Public Health and Preventive Medicine, Monash University, Melbourne, VIC, Australia.

2 Department of Intensive Care, Alfred Hospital, Melbourne, VIC, Australia.

**Correspondence:** alistair.nichol@med.monash.edu.au

### References

1 Hickling KG, Walsh J, Henderson S, Jackson R. Low mortality rate in adult respiratory distress syndrome using low-volume, pressure-

- limited ventilation with permissive hypercapnia: a prospective study. *Crit Care Med* 1994; 22: 1568-78.
- 2 Hickling KG, Henderson SJ, Jackson R. Low mortality associated with low volume pressure limited ventilation with permissive hypercapnia in severe adult respiratory distress syndrome. *Intensive Care Med* 1990; 16: 372-7.
- 3 Amato MB, Barbas CS, Medeiros DM, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med* 1998; 338: 347-54.
- 4 Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med* 2000; 342: 1301-8.
- 5 Terragni PP, Rosboch G, Tealdi A, et al. Tidal hyperinflation during low tidal volume ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2007; 175: 160-6.
- 6 Caironi P, Cressoni M, Chiumello D, et al. Lung opening and closing during ventilation of acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2010; 181: 578-86.
- 7 Hotchkiss JR Jr, Blanch L, Murias G, et al. Effects of decreased respiratory frequency on ventilator-induced lung injury. *Am J Respir Crit Care Med* 2000; 161: 463-8.
- 8 Ranieri VM, Giunta F, Suter PM, Slutsky AS. Mechanical ventilation as a mediator of multisystem organ failure in acute respiratory distress syndrome. *JAMA* 2000; 284: 43-4.
- 9 Kallet RH, Jasmer RM, Luce JM, et al. The treatment of acidosis in acute lung injury with tris-hydroxymethyl aminomethane (THAM). *Am J Respir Crit Care Med* 2000; 161: 1149-53.
- 10 Cooper DJ, Walley KR, Wiggs BR, Russell JA. Bicarbonate does not improve hemodynamics in critically ill patients who have lactic acidosis. A prospective, controlled clinical study. *Ann Intern Med* 1990; 112: 492-8.
- 11 Curley G, Contreras MM, Nichol AD, Higgins BD, Laffey JG. Hypercapnia and acidosis in sepsis: a double-edged sword? *Anesthesiology* 2010; 112: 642-72.
- 12 O’Croinin DF, Nichol AD, Hopkins N, et al. Sustained hypercapnic acidosis during pulmonary infection increases bacterial load and worsens lung injury. *Crit Care Med* 2008; 36: 2128-35.
- 13 O’Croinin DF, Hopkins NO, Moore MM, et al. Hypercapnic acidosis does not modulate the severity of bacterial pneumonia-induced lung injury. *Crit Care Med* 2005; 33: 2606-12.
- 14 Kregenow DA, Rubenfeld GD, Hudson LD, Swenson ER. Hypercapnic acidosis and mortality in acute lung injury. *Crit Care Med* 2006; 34: 1-7.
- 15 Mercat A, Richard JC, Vielle B, et al. Positive end-expiratory pressure setting in adults with acute lung injury and acute respiratory distress syndrome: a randomized controlled trial. *JAMA* 2008; 299: 646-55.
- 16 Meade MO, Cook DJ, Guyatt GH, et al. Ventilation strategy using low tidal volumes, recruitment maneuvers, and high positive end-expiratory pressure for acute lung injury and acute respiratory distress syndrome: a randomized controlled trial. *JAMA* 2008; 299: 637-45.
- 17 Briel M, Meade M, Mercat A, et al. Higher vs lower positive end-expiratory pressure in patients with acute lung injury and acute respiratory distress syndrome: systematic review and meta-analysis. *JAMA* 2010; 303: 865-73.
- 18 Hodgson CL, Tuxen DV, Davies AR, et al. A randomised controlled trial of an open lung strategy with staircase recruitment, titrated PEEP and targeted low airway pressures in patients with acute respiratory distress syndrome. *Crit Care* 2011; 15: R133.
- 19 Cooper DJ, Rosenfeld JV, Murray L, et al. Decompressive craniectomy in diffuse traumatic brain injury. *N Engl J Med* 2011; 364: 1493-502. □